

Sage

## Nose to Toes

**Skin antiseptis, oral cleansing and nasal antiseptis system**

### Competency skill checklist




<b>Name:</b>
<b>Date:</b>

<b>Sage's 2% CHG Cloths</b>			
<b>Gathering product:</b>		<b>Completed</b>	<b>Comments</b>
1.	Inspect warmer for expired product.		
2.	Check settings for appropriate temperature.		
3.	Packages are ready for use when the READY indicator is illuminated.		
4.	Gather the appropriate amount of product based on the facility protocol.		
<b>To open package:</b>		<b>Completed</b>	<b>Comments</b>
1.	Holding the top of the package in one hand, lift the flap on the backside of the package with your other hand.		
2.	Grasp flap at top and pull down to tear flap away and expose foam.		
3.	Hold outside of package to present foam and cloths to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination.		
4.	Transfer contents onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination.		
<b>Warnings:</b>		<b>Completed</b>	<b>Comments</b>
1.	Check with the patient to see if the patient is allergic to chlorhexidine gluconate or any other ingredients in the product. Do not use if they have an allergy.		
2.	Do not use on eyes, ears, and mouth. If contact occurs, rinse with cold water right away and contact a doctor.		

## Nose to Toes

### Skin antisepsis, oral cleansing and nasal antisepsis system Competency skill checklist

Prepping the patient's skin:		Completed	Comments
1.	<p>Follow the protocol chosen by the facility. Indicate prep area(s) on figure below.</p> <p>Fill in protocol here:</p> <div style="text-align: center;">  <p>FRONT      BACK</p> </div>		
2.	Discard any unused cloths.		

### 3M's Skin and Nasal Antiseptic

Application instructions:	Completed	Comments
Open package and remove bottle swabs.		
Unscrew cap by turning cap counterclockwise.		
Use a tissue to clean the inside of both nostrils including the inside tip of nostril. Discard.		
Tilting the bottle slightly, dip one swab into solution and stir vigorously for 10 seconds. Withdraw the swab slowly to avoid wiping solution off during removal.		
Insert swab comfortably into one nostril and rotate for 15 seconds covering all surfaces. Then focus on the inside tip of nostril and rotate for an additional 15 seconds. (swab 1). Repeat for swab 2, 3, and 4.		
Discard product.		

Sage

**Nose to Toes**

**Skin antisepsis, oral cleansing and nasal antisepsis system**  
**Competency skill checklist**

<b>Oral Care</b>		
<b>Using the single dose bottle of CHG .12% oral infection:</b>	<b>Completed</b>	<b>Comments</b>
Allow patient to rinse with the antiseptic and spit in basin.		
Discard after use.		

\_\_\_\_\_  
**Employee signature**

\_\_\_\_\_  
**Observer signature**