

Sage

Skin and nasal antiseptis

Competency skill checklist




Name:
Date:

Sage's 2% CHG Cloths			
Gathering product:		Completed	Comments
1.	Inspect warmer for expired product.		
2.	Check settings for appropriate temperature.		
3.	Packages are ready for use when the READY indicator is illuminated.		
4.	Gather the appropriate amount of product based on the facility protocol.		
To open package:		Completed	Comments
1.	Holding the top of the package in one hand, lift the flap on the backside of the package with your other hand.		
2.	Grasp flap at top and pull down to tear flap away and expose foam.		
3.	Hold outside of package to present foam and cloths to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination.		
4.	Transfer contents onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination.		
Warnings:		Completed	Comments
1.	Check with the patient to see if the patient is allergic to chlorhexidine gluconate or any other ingredients in the product. Do not use if they have an allergy.		
2.	Do not use on eyes, ears, and mouth. If contact occurs, rinse with cold water right away and contact a doctor.		
Prepping the patient's skin:		Completed	Comments

Sage

Skin and nasal antiseptis

Competency skill checklist

<p>1.</p>	<p>Follow the protocol chosen by the facility. Indicate prep area(s) on figure below.</p> <p>Fill in protocol here:</p> <div style="text-align: center;">  </div>		
<p>2.</p>	<p>Discard any unused cloths.</p>		

3M's Skin and Nasal Antiseptic

Application instructions:	Completed	Comments
Open package and remove bottle swabs.		
Unscrew cap by turning cap counterclockwise.		
Use a tissue to clean the inside of both nostrils including the inside tip of nostril. Discard.		
Tilting the bottle slightly, dip one swab into solution and stir vigorously for 10 seconds. Withdraw the swab slowly to avoid wiping solution off during removal.		
Insert swab comfortably into one nostril and rotate for 15 seconds covering all surfaces. Then focus on the inside tip of nostril and rotate for an additional 15 seconds. (swab 1). Repeat for swab 2, 3, and 4.		
Discard of product.		

Employee signature

Observer signature